



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Solution by PROFESSOR F. L. GRIFFIN, Williams College.

However the ellipse is placed relative to the co-ordinate axes, its area is

$$S = \int_{x_1}^{x_2} (y'' - y') dx,$$

where x_1 and x_2 [$x_1 < x_2$] are the extreme abscissas taken in the curve, and y' and y'' [$y' \leq y''$] are the two ordinates corresponding to any one value of x . Solved for y , the given equation becomes

$$by = -(hx + f) \pm \sqrt{[f^2 - bc + 2(hf - bg)x + (h^2 - ab)x^2]}.$$

Now let

$$f^2 - bc = b^2 A, \quad fh - bg = b^2 B, \quad ab - h^2 = b^2 C,$$

where for an ellipse $C > 0$. Then the difference of the two ordinates becomes

$$y'' - y' = 2\sqrt{[A + 2Bx - Cx^2]}.$$

Hence, integrating,

$$S = 2 \left[\frac{Cx - B}{2C} \sqrt{[A + 2Bx - Cx^2]} + \frac{B^2 + AC}{2C^{\frac{3}{2}}} \sin^{-1} \frac{Cx - B}{\sqrt{[B^2 + AC]}} \right]_{x_1}^{x_2}$$

Now the extreme abscissas make $y' = y''$, or $A + 2Bx - Cx^2 = 0$; whence

$$Cx_2 = B + \sqrt{[B^2 + AC]} \quad \text{and} \quad Cx_1 = B - \sqrt{[B^2 + AC]}.$$

Substituting these values,

$$S = \frac{B^2 + AC}{C^{\frac{3}{2}}} [\sin^{-1} 1 - \sin^{-1}(-1)] = \frac{\pi}{b} \frac{[(fh - bg)^2 + (ab - h^2)(f^2 - bc)]}{(ab - h^2)^{\frac{3}{2}}},$$

which immediately reduces to the formula proposed.

Also solved by G. B. M. Zerr and J. Scheffer.

MECHANICS.

240. Proposed by S. A. COREY, Hiteman, Iowa.

A perfectly flexible wire rope weighing one pound per foot is suspended from the tops of two vertical supports 300 feet apart, one support being 30 feet higher than the other. One end of the rope is fastened to the top of the higher support, while 600 feet of the rope hangs vertically from the top of the lower support. Assuming that the rope is free to slide over the top of the lower support without friction, find the lowest point of

that portion of the rope which is suspended between the supports. Also find the amount of work which must be performed in raising the lowest point to make it coincide with the top of the lower support by exerting a pull on the free end of the rope.

[No solution of this problem has been received.]

240. Proposed by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.

A simple beam length $2a$, supported at both ends, is loaded in the form of a parabola, height of vertex b . Find deflection at center due to this load.

Solution by the PROPOSER.

Let AB be the beam, ACD the parabola, $CD=b$, $AD=DB=a$. Take E any point on AB , draw EF perpendicular to AB . Let $AE=x$, and also z be the distance of the center of gravity of the area AEF from EF . Then $\frac{2}{3}ab$ =total load. $(x-a)^2 + (a^2/b)(y-b)=0$ is the equation to the parabola, with A as origin.

$$\text{Then } (x-z) = \frac{\int \int x dx dy}{\int \int dx dy} = \frac{\int \int x dx dy}{A}.$$

$$\therefore A(x-z) = \int_0^x xy dx = \frac{b}{a^2} \int_0^x (2ax^2 - x^3) dx = \frac{bx^3}{12a^2} (8a-3x).$$

$$A = \int_0^x y dx = \frac{bx^2}{3a^2} (3a-x).$$

$$\therefore x-z = \frac{8ax-3x^2}{4(3a-x)} \text{ and } z = \frac{4ax-x^2}{4(3a-x)}.$$

Taking moments about A we get,

$$EI \frac{d^2 y}{dx^2} = \frac{2}{3}abx - Az = \frac{2}{3}abx - \frac{bx^3}{3a} + \frac{bx^4}{12a^2} = M.$$

$$EI \frac{dy}{dx} = \frac{1}{3}abx^2 - \frac{bx^4}{12a} + \frac{bx^5}{60a^2} + C.$$

When $x=a$, $dy/dx=0$, $C = -\frac{4}{15}a^3b$.

$$\therefore EI \frac{dy}{dx} = \frac{1}{3}abx^2 - \frac{bx^4}{12a} + \frac{bx^5}{60a^2} - \frac{4}{15}a^3b,$$

$$EI y = \frac{1}{9}abx^3 - \frac{bx^5}{60a} + \frac{bx^6}{360a^2} - \frac{4}{15}a^3bx = -\frac{61a^4b}{360} \text{ when } x=a.$$